INFANT AND CHILDREN'S ACETAMINOPHEN & IBUPROFEN SHORTAGE OPTIONS

With the recent shortage of infant and children's liquid and chewable tablets of acetaminophen (Tylenol[®]) and ibuprofen (Advil[®] or Motrin[®]), parents can consider splitting and/or crushing adult tablets and mixing with a small amount of juice, apple sauce or pudding for use. Any extended-release products that are used every 8 hours or more (e.g. Tylenol Arthritis[®]) should NOT be used in this way.

Acetaminophen for infants/children <12y of age:

Adult tablets are available over-the-counter as REGULAR 325 mg tablets OR as EXTRA STRENGTH 500mg tablets and can be crushed. **** DO NOT USE EXTENDED-RELEASE PRODUCTS OF ACETAMINOPHEN****

Suppositories are available over-the-counter as 120 mg, 160 mg, 325 mg and 650 mg suppositories.

The following table can be used for dosing. Each dose should be given every 4 hours; maximum of 5 doses a day.

Kg	Lbs	Age	Usual Dose	Suggested oral tablet option	Proportion of Tablet	Suggested rectal option
2.7 to 5.3	6 to 11	0 to 3 mo	40 mg	40.62 mg (1/8 tab of 325 mg tab)		40 mg (1/4 of 160 mg supp)
5.4 to 8.1	12 to 17	4 to 11 mo	80 mg	81.25 mg (1/4 tab of 325 mg tab)		80 mg (1/2 of 160 mg supp)
8.2 to 10.8	18 to 23	1 to 2 y	120 mg	125 mg (1/4 tab of 500 mg tab) OR 81.25 mg (1/4 tab of 325 mg tab) for cautious dosing	(325mg or 500 mg)	120 mg supp
10.9 to 16.3	24 to 35	2 to 3 y	160 mg	162.5 mg (1/2 tab of 325 mg tab)		160 mg supp
16.4 to 21.7	36 to 47	4 to 5 y	240 mg	250 mg (1/2 tab of 500 mg tab) OR 243 mg (3/4 tab of 325 mg tab)	500mg or 325mg	240 mg (2 x 120mg supp)
21.8 to 27.2	48 to 59	6 to 8 y	320 mg to 325 mg	325 mg tab		325 mg supp
27.3 to 32.6	60 to 71	9 to 10 y	325 mg to 400 mg	325 mg tab		325 mg supp
32.7 to 43.2	72 to 95	11 y	480 mg to 500 mg	500 mg tab		650 mg supp

Adapted and Reviewed by: Brenda Chang, Doret Cheng & Jon Hunchuck St. Michael's Academic Family Health Team Pharmacists

INFANT AND CHILDREN'S ACETAMINOPHEN & IBUPROFEN SHORTAGE OPTIONS

Ibuprofen for infants/children 6 mo to <12y of age:

Adult tablets are available over-the-counter as 200mg, 300mg, and 400mg tablets and can be crushed. **** DO NOT USE ANY EXTENDED-RELEASE PRODUCTS OF IBUPROFEN****

The following table can be used for dosing. Each dose should be given every 6 to 8 hours; maximum of 4 doses per day.

Кg	Lbs	Age	Usual Dose	Suggested oral tablet option	Proportion of Tablet
5.4 to 8.1	12 to 17	6 to 11 mo	50 mg	50 mg (1/4 tab of <mark>200 mg</mark> tab)	
8.2 to 10.8	18 to 23	1 to 2 y	75 to 80 mg	75 mg (1/4 tab of 300 mg tab) OR 50 mg (1/4 tab of 200 mg tab) for cautious dosing	300mg or 200mg
10.9 to 16.3	24 to 35	2 to 3 y	100 mg	100 mg (1/2 tab of <mark>200 mg</mark> tab)	
16.4 to 21.7	36 to 47	4 to 5 y	150 mg	150 mg (1/2 tab of 300 mg tab) OR 150 mg (3/4 tab of 200mg tab)	300mg Or 200mg
21.8 to 27.2	48 to 59	6 to 8 y	200 mg	200 mg tab	\bullet
27.3 to 32.6	60 to 71	9 to 10 y	200 mg to 250 mg	200 mg tab	\bullet
32.7 to 43.2	72 to 95	11 y	300 mg	300 mg tab OR 300mg (1 and ½ of 200mg tab)	300mg Or 200mg

References:

1. Acetaminophen. In: Pediatric & Neonatal Lexi-Drugs Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.: 2022 [updated 12/11/2022; cited 14/11/2022]. Available from: http://online.lexi.com. Subscription required to view.

2. Ibuprofen. In: Pediatric & Neonatal Lexi-Drugs Online Database [database on the Internet]. Hudson (OH): Lexicomp Inc.: 2022 [updated 12/11/2022; cited 14/11/2022]. Available from: http://online.lexi.com. Subscription required to view.

Adapted and Reviewed by: Brenda Chang, Doret Cheng & Jon Hunchuck St. Michael's Academic Family Health Team Pharmacists